

Generating all subsets of a finite set with disjoint unions

David Ellis*, Benny Sudakov†

May 2011

Abstract

If X is an n -element set, we call a family $\mathcal{G} \subset \mathcal{P}X$ a k -generator for X if every $x \subset X$ can be expressed as a union of at most k disjoint sets in $\mathcal{G}$. Frein, Lévêque and Sebő [10] conjectured that for $n > 2k$, the smallest k -generators for X are obtained by taking a partition of X into classes of sizes as equal as possible, and taking the union of the power-sets of the classes. We prove this conjecture for all sufficiently large n when $k = 2$, and for n a sufficiently large multiple of k when $k \geq 3$.

Keywords: generator, disjoint unions.
2000 MSC: 05Dxx.

1 Introduction

Let X be an n -element set, and let $\mathcal{P}X$ denote the set of all subsets of X . We call a family $\mathcal{G} \subset \mathcal{P}X$ a k -generator for X if every $x \subset X$ can be expressed as a union of at most k disjoint sets in $\mathcal{G}$. For example, let $(V_i)_{i=1}^k$ be a partition of X into k classes of sizes as equal as possible; then

$$\mathcal{F}_{n,k} := \bigcup_{i=1}^k \mathcal{P}(V_i) \setminus \{\emptyset\}$$

*St John's College, Cambridge, CB2 1TP, UK.

†Department of Mathematics, UCLA, Los Angeles, CA 90095, USA. Research supported in part by NSF CAREER award DMS-0812005 and by a USA-Israeli BSF grant.

is a k -generator for X . We call a k -generator of this form *canonical*. If $n = qk + r$, where $0 \leq r < k$, then

$$|\mathcal{F}_{n,k}| = (k - r)(2^q - 1) + r(2^{q+1} - 1) = (k + r)2^q - k.$$

Frein, L  v  que and Seb   [10] conjectured that for any $k \leq n$, this is the smallest possible size of a k -generator for X .

Conjecture 1 (Frein, L  v  que, Seb  ). *If X is an n -element set, $k \leq n$, and $\mathcal{G} \subset \mathcal{P}X$ is a k -generator for X , then $|\mathcal{G}| \geq |\mathcal{F}_{n,k}|$. If $n > 2k$, equality holds only if $\mathcal{G}$ is a canonical k -generator for X .*

They proved this for $k \leq n \leq 3k$, but their methods do not seem to work for larger n .

For $k = 2$, Conjecture 1 is a weakening of a conjecture of Erd  s. We call a family $\mathcal{G} \subset \mathcal{P}X$ a k -base for X if every $x \subset X$ can be expressed as a union of at most k (not necessarily disjoint) sets in $\mathcal{G}$. Erd  s (see [11]) made the following

Conjecture 2 (Erd  s). *If X is an n -element set, and $\mathcal{G} \subset \mathcal{P}X$ is a 2-base for X , then $|\mathcal{G}| \geq |\mathcal{F}_{n,2}|$.*

In fact, Frein, L  v  que and Seb   [10] made the analogous conjecture for all k .

Conjecture 3 (Frein, L  v  que, Seb  ). *If X is an n -element set, $k \leq n$, and $\mathcal{G} \subset \mathcal{P}X$ is a k -base for X , then $|\mathcal{G}| \geq |\mathcal{F}_{n,k}|$. If $n > 2k$, equality holds only if $\mathcal{G}$ is a canonical k -generator for X .*

Again, they were able to prove this for $k \leq n \leq 3k$.

In this paper, we study k -generators when n is large compared to k . Our main results are as follows.

Theorem 4. *If n is sufficiently large, X is an n -element set, and $\mathcal{G} \subset \mathcal{P}X$ is a 2-generator for X , then $|\mathcal{G}| \geq |\mathcal{F}_{n,2}|$. Equality holds only if $\mathcal{G}$ is of the form $\mathcal{F}_{n,2}$.*

Theorem 5. *If $k \in \mathbb{N}$, n is a sufficiently large multiple of k , X is an n -element set, and $\mathcal{G}$ is a k -generator for X , then $|\mathcal{G}| \geq |\mathcal{F}_{n,k}|$. Equality holds only if $\mathcal{G}$ is of the form $\mathcal{F}_{n,k}$.*

In other words, we prove Conjecture 1 for all sufficiently large n when $k = 2$, and for n a sufficiently large multiple of k when $k \geq 3$. We use some ideas of Alon and Frankl [1], and also techniques of the first author from [5], in which asymptotic results were obtained.

As noted in [10], if $\mathcal{G} \subset \mathcal{P}X$ is a k -generator (or even a k -base) for X , then the number of ways of choosing at most k sets from $\mathcal{G}$ is clearly at least the number of subsets of X . Therefore $|\mathcal{G}|^k \geq 2^n$, which immediately gives

$$|\mathcal{G}| \geq 2^{n/k}.$$

Moreover, if $|\mathcal{G}| = m$, then

$$\sum_{i=0}^k \binom{m}{i} \geq 2^n. \quad (1)$$

Crudely, we have

$$\sum_{i=0}^{k-1} \binom{m}{i} \leq 2m^{k-1},$$

so

$$\sum_{i=0}^k \binom{m}{i} \leq \binom{m}{k} + 2m^{k-1}.$$

Hence, if k is fixed, then

$$(1 + O(1/m)) \binom{m}{k} \geq 2^n,$$

so

$$|\mathcal{G}| \geq (k!)^{1/k} 2^{n/k} (1 - o(1)). \quad (2)$$

Observe that if $n = qk + r$, where $0 \leq r < k$, then

$$|\mathcal{F}_{n,k}| = (k+r)2^q - k < (k+r)2^q = k2^{n/k}(1+r/k)2^{-r/k} < c_0 k 2^{n/k}, \quad (3)$$

where

$$c_0 := \frac{2}{2^{1/\log 2} \log 2} = 1.061 \text{ (to 3 d.p.)}.$$

Now for some preliminaries. We use the following standard notation. For $n \in \mathbb{N}$, $[n]$ will denote the set $\{1, 2, \dots, n\}$. If x and y are disjoint sets, we will sometimes write their union as $x \sqcup y$, rather than $x \cup y$, to emphasize the fact that the sets are disjoint.

If $k \in \mathbb{N}$, and G is a graph, $K_k(G)$ will denote the number of k -cliques in G . Let $T_s(n)$ denote the s -partite Turán graph (the complete s -partite graph on n vertices with parts of sizes as equal as possible), and let $t_s(n) = e(T_s(n))$. For $l \in \mathbb{N}$, C_l will denote the cycle of length l .

If F is a (labelled) graph on f vertices, with vertex-set $\{v_1, \dots, v_f\}$ say, and $\mathbf{t} = (t_1, \dots, t_f) \in \mathbb{N}^f$, we define the $\mathbf{t}$ -blow-up of F , $F \otimes \mathbf{t}$, to be the

graph obtained by replacing v_i with an independent set V_i of size t_i , and joining each vertex of V_i to each vertex of V_j whenever $v_i v_j$ is an edge of F . With slight abuse of notation, we will write $F \otimes t$ for the symmetric blow-up $F \otimes (t, \dots, t)$.

If F and G are graphs, we write $c_F(G)$ for the number of injective graph homomorphisms from F to G , meaning injections from $V(F)$ to $V(G)$ which take edges of F to edges of G . The *density of F in G* is defined to be

$$d_F(G) = \frac{c_F(G)}{|G|(|G| - 1) \cdots (|G| - |F| + 1)},$$

i.e. the probability that a uniform random injective map from $V(F)$ to $V(G)$ is a graph homomorphism from F to G . Hence, when $F = K_k$, the density of K_k 's in an n -vertex graph G is simply $K_k(G)/\binom{n}{k}$.

Although we will be interested in the density $d_F(G)$, it will sometimes be more convenient to work with the following closely related quantity, which behaves very nicely when we take blow-ups. We write $\text{Hom}_F(G)$ for the number of homomorphisms from F to G , and we define the *homomorphism density of F in G* to be

$$h_F(G) = \frac{\text{Hom}_F(G)}{|G|^{|F|}},$$

i.e. the probability that a uniform random map from $V(F)$ to $V(G)$ is a graph homomorphism from F to G .

Observe that if F is a graph on f vertices, and G is a graph on n vertices, then the number of homomorphisms from F to G which are not injections is clearly at most

$$\binom{f}{2} n^{f-1}.$$

Hence,

$$d_G(F) \geq \frac{h_G(F)n^f - \binom{f}{2}n^{f-1}}{n(n-1) \cdots (n-f+1)} \geq h_G(F) - O(1/n), \quad (4)$$

if f is fixed. In the other direction,

$$d_F(G) \leq \frac{n^f}{n(n-1) \cdots (n-f+1)} h_F(G) \leq (1 + O(1/n)) h_F(G) \quad (5)$$

if f is fixed. Hence, when working inside large graphs, we can pass freely between the density of a fixed graph F and its homomorphism density, with an ‘error’ of only $O(1/n)$.

Finally, we will make frequent use of the AM/GM inequality:

Theorem 6. *If $x_1, \dots, x_n \geq 0$, then*

$$\left(\prod_{i=1}^n x_i \right)^{1/n} \leq \frac{1}{n} \sum_{i=1}^n x_i.$$

2 The case $k \mid n$ via extremal graph theory.

For n a sufficiently large multiple of k , it turns out to be possible to prove Conjecture 1 using stability versions of Turán-type results. We will prove the following

Theorem 5. *If $k \in \mathbb{N}$, n is a sufficiently large multiple of k , X is an n -element set, and $\mathcal{G}$ is a k -generator for X , then $|\mathcal{G}| \geq |\mathcal{F}_{n,k}|$. Equality holds only if $\mathcal{G}$ is of the form $\mathcal{F}_{n,k}$.*

We need a few more definitions. Let H denote the graph with vertex-set $\mathcal{P}X$, where we join two subsets $x, y \subset X$ if they are disjoint. With slight abuse of terminology, we call H the ‘Kneser’ graph on $\mathcal{P}X$ (although this usually means the analogous graph on $X^{(r)}$). If $\mathcal{F}, \mathcal{G} \subset \mathcal{P}X$, we say that $\mathcal{G}$ *k-generates* $\mathcal{F}$ if every set in $\mathcal{F}$ is a disjoint union of at most k sets in $\mathcal{G}$.

The main steps of the proof: First, we will show that for any $\mathcal{A} \subset \mathcal{P}X$ with $|\mathcal{A}| \geq \Omega(2^{n/k})$, the density of K_{k+1} ’s in the induced subgraph $H[\mathcal{A}]$ is $o(1)$.

Secondly, we will observe that if n is a sufficiently large multiple of k , and $\mathcal{G} \subset \mathcal{P}X$ has size close to $|\mathcal{F}_{n,k}|$ and k -generates almost all subsets of X , then $K_k(H[\mathcal{G}])$ is very close to $K_k(T_k(|\mathcal{G}|))$, the number of K_k ’s in the k -partite Turán graph on $|\mathcal{G}|$ vertices.

We will then prove that if G is any graph with small K_{k+1} -density, and with $K_k(G)$ close to $K_k(T_k(|G|))$, then G can be made k -partite by removing a small number of edges. This can be seen as a (strengthened) variant of the Simonovits Stability Theorem [9], which states that any K_{k+1} -free graph G with $e(G)$ close to the maximum $e(T_k(|G|))$, can be made k -partite by removing a small number of edges.

This will enable us to conclude that $H[\mathcal{G}]$ can be made k -partite by the removal of a small number of edges, and therefore the structure of $H[\mathcal{G}]$ is close to that of the Turán graph $T_k(|\mathcal{G}|)$. This in turn will enable us to show that the structure of $\mathcal{G}$ is close to that of a canonical k -generator $\mathcal{F}_{n,k}$ (Proposition 9).

Finally, we will use a perturbation argument to show that if n is sufficiently large, and $|\mathcal{G}| \leq |\mathcal{F}_{n,k}|$, then $\mathcal{G} = \mathcal{F}_{n,k}$, completing the proof.

In fact, we will first show that if $\mathcal{A} \subset \mathcal{P}X$ with $|\mathcal{A}| \geq \Omega(2^{n/k})$, then the homomorphism density of $K_{k+1} \otimes t$ in $H[\mathcal{A}]$ is $o(1)$, provided t is sufficiently large depending on k . Hence, we will need the following (relatively well-known) lemma relating the homomorphism density of a graph to that of its blow-up.

Lemma 7. *Let F be a graph on f vertices, let $\mathbf{t} = (t_1, t_2, \dots, t_f) \in \mathbb{N}^f$, and let $F \otimes \mathbf{t}$ denote the $\mathbf{t}$ -blow-up of F . If the homomorphism density of F in G is p , then the homomorphism density of $F \otimes \mathbf{t}$ in G is at least $p^{t_1 t_2 \dots t_f}$.*

Proof. This is a simple convexity argument, essentially that of [9]. It will suffice to prove the statement of the lemma when $\mathbf{t} = (1, \dots, 1, r)$ for some $r \in \mathbb{N}$. We think of F as a (labelled) graph on vertex set $[f] = \{1, 2, \dots, f\}$, and G as a (labelled) graph on vertex set $[n]$. Define the function $\chi : [n]^f \rightarrow \{0, 1\}$ by

$$\chi(v_1, \dots, v_f) = \begin{cases} 1 & \text{if } i \mapsto v_i \text{ is a homomorphism from } F \text{ to } G, \\ 0 & \text{otherwise.} \end{cases}$$

Then we have

$$h_F(G) = \frac{1}{n^f} \sum_{(v_1, \dots, v_f) \in [n]^f} \chi(v_1, \dots, v_f) = p.$$

The homomorphism density $h_{F \otimes (1, \dots, 1, r)}(G)$ of $F \otimes (1, \dots, 1, r)$ in G is:

$$\begin{aligned} h_{F \otimes (1, \dots, 1, r)}(G) &= \frac{1}{n^{f-1+r}} \sum_{(v_1, \dots, v_{f-1}, v_f^{(1)}, v_f^{(2)}, \dots, v_f^{(r)}) \in [n]^{f-1+r}} \prod_{i=1}^r \chi(v_1, \dots, v_{f-1}, v_f^{(i)}) \\ &= \frac{1}{n^{f-1}} \sum_{(v_1, \dots, v_{f-1}) \in [n]^{f-1}} \left(\frac{1}{n} \sum_{v_f \in [n]} \chi(v_1, \dots, v_{f-1}, v_f) \right)^r \\ &\geq \left(\frac{1}{n^{f-1}} \sum_{(v_1, \dots, v_{f-1}) \in [n]^{f-1}} \left(\frac{1}{n} \sum_{v_f \in [n]} \chi(v_1, \dots, v_{f-1}, v_f) \right) \right)^r \\ &= \left(\frac{1}{n^f} \sum_{(v_1, \dots, v_{f-1}, v_f) \in [n]^f} \chi(v_1, \dots, v_{f-1}, v_f) \right)^r \\ &= p^r. \end{aligned}$$

Here, the inequality follows from applying Jensen's Inequality to the convex function $x \mapsto x^r$. This proves the lemma for $\mathbf{t} = (1, \dots, 1, r)$. By

symmetry, the statement of the lemma holds for all vectors of the form $(1, \dots, 1, r, 1, \dots, 1)$. Clearly, we may obtain $F \otimes \mathbf{t}$ from F by a sequence of blow-ups by these vectors, proving the lemma. $\square$

The following lemma (a rephrasing of Lemma 4.2 in Alon and Frankl [1]) gives an upper bound on the homomorphism density of $K_{k+1} \otimes t$ in large induced subgraphs of the Kneser graph H .

Lemma 8. *If $\mathcal{A} \subset \mathcal{P}X$ with $|\mathcal{A}| = m = 2^{(\delta+1/(k+1))n}$, then*

$$h_{K_{k+1} \otimes t}(H[\mathcal{A}]) \leq (k+1)2^{-n(\delta t-1)}.$$

Proof. We follow the proof of Alon and Frankl cited above. Choose $(k+1)t$ members of $\mathcal{A}$ uniformly at random with replacement, $(A_i^{(j)})_{1 \leq i \leq k+1, 1 \leq j \leq t}$. The homomorphism density of $K_{k+1} \otimes t$ in $H[\mathcal{A}]$ is precisely the probability that the unions

$$U_i = \bigcup_{j=1}^t A_i^{(j)}$$

are pairwise disjoint. If this event occurs, then $|U_i| \leq n/(k+1)$ for some i . For each $i \in [k]$, we have

$$\begin{aligned} \Pr\{|U_i| \leq n/(k+1)\} &= \Pr\left(\bigcup_{S \subset X: |S| \leq n/(k+1)} \left(\bigcap_{j=1}^t \{A_i^{(j)} \subset S\}\right)\right) \\ &\leq \sum_{|S| \leq n/(k+1)} \Pr\left(\bigcap_{j=1}^t \{A_i^{(j)} \subset S\}\right) \\ &= \sum_{|S| \leq n/(k+1)} (2^{|S|}/m)^t \\ &\leq 2^n (2^{n/(k+1)}/m)^t \\ &= 2^{-n(\delta t-1)}. \end{aligned}$$

Hence,

$$\Pr\left(\bigcup_{i=1}^k \{|U_i| \leq n/(k+1)\}\right) \leq \sum_{i=1}^k \Pr\{|U_i| \leq n/(k+1)\} \leq (k+1)2^{-n(\delta t-1)}.$$

Therefore,

$$h_{K_{k+1} \otimes t}(H[\mathcal{A}]) \leq (k+1)2^{-n(\delta t-1)},$$

as required. $\square$

From the trivial bound above, any k -generator $\mathcal{G}$ has $|\mathcal{G}| \geq 2^{n/k}$, so $\delta \geq 1/(k(k+1))$, and therefore, choosing $t = t_k := 2k(k+1)$, we see that

$$h_{K_{k+1} \otimes t_k}(H[\mathcal{G}]) \leq (k+1)2^{-n}.$$

Hence, by Lemma 7,

$$h_{K_{k+1}}(H[\mathcal{G}]) \leq O_k(2^{-n/t_k^k}).$$

Therefore, by (5),

$$d_{K_{k+1}}(H[\mathcal{G}]) \leq O_k(2^{-n/t_k^k}) \leq 2^{-a_k n} \quad (6)$$

provided n is sufficiently large depending on k , where $a_k > 0$ depends only on k .

Assume now that n is a multiple of k , so that $|\mathcal{F}_{n,k}| = k2^{n/k} - k$. We will prove the following ‘stability’ result.

Proposition 9. *Let $k \in \mathbb{N}$ be fixed. If n is a multiple of k , and $\mathcal{G} \subset \mathcal{P}X$ has $|\mathcal{G}| \leq (1+\eta)|\mathcal{F}_{n,k}|$ and k -generates at least $(1-\epsilon)2^n$ subsets of X , then there exists an equipartition $(S_i)_{i=1}^k$ of X such that*

$$|\mathcal{G} \cap (\cup_{i=1}^k \mathcal{P}S_i)| \geq (1 - C_k \epsilon^{1/k} - D_k \eta^{1/k} - 2^{-\xi_k n})|\mathcal{F}_{n,k}|,$$

where $C_k, D_k, \xi_k > 0$ depend only on k .

We first collect some results used in the proof. We will need the following theorem of Erdős [7].

Theorem 10 (Erdős). *If $r \leq k$, and G is a K_{k+1} -free graph on n vertices, then*

$$K_r(G) \leq K_r(T_k(n)).$$

We will also need the following well-known lemma, which states that a dense k -partite graph has an induced subgraph with high minimum degree.

Lemma 11. *Let G be an n -vertex, k -partite graph with*

$$e(G) \geq (1 - 1/k - \delta)n^2/2.$$

Then there exists an induced subgraph $G' \subseteq G$ with $|G'| = n' \geq (1 - \sqrt{\delta})n$ and minimum degree $\delta(G') \geq (1 - 1/k - \sqrt{\delta})(n' - 1)$.

Proof. We perform the following algorithm to produce G' . Let $G_1 = G$. Suppose that at stage i , we have a graph G_i on $n - i + 1$ vertices. If there is a vertex v of G_i with $d(v) < (1 - 1/k - \eta)(n - i)$, let $G_{i+1} = G_i - v$; otherwise, stop and set $G' = G_i$. Suppose the process terminates after $j = \alpha n$ steps. Then we have removed at most

$$(1 - 1/k - \eta) \sum_{i=1}^j (n - i) = (1 - 1/k - \eta) \left(\binom{n}{2} - \binom{n-j}{2} \right)$$

edges, and the remaining graph has at most

$$\binom{k}{2} \left(\frac{n-j}{k} \right)^2 = (1 - \alpha)^2 (1 - 1/k) n^2 / 2$$

edges. But our original graph had at least

$$(1 - 1/k - \delta) n^2 / 2$$

edges, and therefore

$$(1 - 1/k - \eta)(1 - (1 - \alpha)^2) n^2 / 2 + (1 - \alpha)^2 (1 - 1/k) n^2 / 2 \geq (1 - 1/k - \delta) n^2 / 2,$$

so

$$\eta(1 - \alpha)^2 \geq \eta - \delta.$$

Choosing $\eta = \sqrt{\delta}$, we obtain

$$\eta(1 - \alpha)^2 \geq \eta(1 - \eta),$$

and therefore

$$(1 - \alpha)^2 \geq 1 - \eta,$$

so

$$\alpha \leq 1 - (1 - \eta)^{1/2} \leq \eta.$$

Hence, our induced subgraph G' has order

$$|G'| = n' \geq (1 - \sqrt{\delta})n,$$

and minimum degree

$$\delta(G') \geq (1 - 1/k - \sqrt{\delta})(n' - 1).$$

□

We will also need Shearer's Entropy Lemma.

Lemma 12 (Shearer's Entropy Lemma, [4]). *Let S be a finite set, and let $\mathcal{A}$ be an r -cover of S , meaning a collection of subsets of S such that every element of S is contained in at least r sets in $\mathcal{A}$. Let $\mathcal{F}$ be a collection of subsets of S . For $A \subset S$, let $\mathcal{F}_A = \{F \cap A : F \in \mathcal{F}\}$ denote the projection of $\mathcal{F}$ onto the set A . Then*

$$|\mathcal{F}|^r \leq \prod_{A \in \mathcal{A}} |\mathcal{F}_A|.$$

In addition, we require two 'stability' versions of Turán-type results in extremal graph theory. The first states that a graph with a very small K_{k+1} -density cannot have K_r -density much higher than the k -partite Turán graph on the same number of vertices, for any $r \leq k$.

Lemma 13. *Let $r \leq k$ be integers. Then there exist $C, D > 0$ such that for any $\alpha \geq 0$, any n -vertex graph G with K_{k+1} -density at most α has K_r -density at most*

$$\frac{k(k-1) \cdots (k-r+1)}{k^r} (1 + C\alpha^{1/(k+2)} + D/n).$$

Proof. We use a straightforward sampling argument. Let G be as in the statement of the lemma. Let $\zeta \binom{n}{l}$ be the number of l -subsets $U \subset V(G)$ such that $G[U]$ contains a copy of K_{k+1} , so that ζ is simply the probability that a uniform random l -subset of $V(G)$ contains a K_{k+1} . Simple counting (or the union bound) gives

$$\zeta \leq \binom{l}{k+1} \alpha.$$

By Theorem 10, each K_{k+1} -free $G[U]$ contains at most

$$\binom{k}{r} \left(\frac{l}{k} \right)^r$$

K_r 's. Therefore, the density of K_r 's in each such $G[U]$ satisfies

$$\begin{aligned} d_{K_r}(G[U]) &\leq \frac{k(k-1) \cdots (k-r+1)}{k^r} \frac{l^r}{l(l-1) \cdots (l-r+1)} \\ &\leq \frac{k(k-1) \cdots (k-r+1)}{k^r} (1 + O(1/l)). \end{aligned} \tag{7}$$

Note that one can choose a random r -set in graph G by first choosing a random l -set U , and then choosing a random r -subset of U . The density of

K_r 's in G is simply the probability that a uniform random r -subset of $V(G)$ induces a K_r , and therefore

$$d_{K_r}(G) = \mathbb{E}_U[d_{K_k}(G[U])],$$

where the expectation is taken over a uniform random choice of U . If U is K_{k+1} -free, which happens with probability $1 - \zeta$, we use the upper bound (7); if U contains a K_{k+1} , which happens with probability ζ , we use the trivial bound $d_{K_k}(G[U]) \leq 1$. We see that the density of K_r 's in G satisfies:

$$\begin{aligned} d_{K_r}(G) &\leq (1 - \zeta) \frac{k(k-1) \cdots (k-r+1)}{k^r} (1 + O(1/l)) + \zeta \\ &\leq \frac{k(k-1) \cdots (k-r+1)}{k^r} + O(1/l) + \binom{l}{k+1} \alpha \\ &\leq \frac{k(k-1) \cdots (k-r+1)}{k^r} + O(1/l) + l^{k+1} \alpha. \end{aligned}$$

Choosing $l = \min\{\lfloor \alpha^{-1/(k+2)} \rfloor, n\}$ proves the lemma. $\square$

The second result states that an n -vertex graph with a small K_{k+1} -density, a K_k -density not too much less than that of $T_k(n)$, and a K_{k-1} -density not too much more than that of $T_k(n)$, can be made into a k -partite graph by the removal of only a small number of edges.

Theorem 14. *Let G be an n -vertex graph with K_{k+1} -density at most α , K_{k-1} -density at most*

$$(1 + \beta) \frac{k!}{k^{k-1}},$$

and K_k -density at least

$$(1 - \gamma) \frac{k!}{k^k},$$

where $\gamma \leq 1/2$. Then G can be made into a k -partite graph G_0 by removing at most

$$\left(2\beta + 2\gamma + \frac{8k^{k+1}(k+1)}{k!} \sqrt{\alpha} + 2k/n \right) \binom{n}{2}$$

edges, which removes at most

$$\left(2\beta + 2\gamma + \frac{8k^{k+1}(k+1)}{k!} \sqrt{\alpha} + 2k/n \right) \binom{k}{2} \binom{n}{k}$$

K_k 's.

Proof. If $k \in \mathbb{N}$, and G is a graph, let

$$\mathcal{K}_k(G) = \{S \in V(G)^{(k)} : G[S] \text{ is a clique}\}$$

denote the set of all k -sets that induce a clique in G . If $S \subset V(G)$, let $N(S)$ denote the set of vertices of G joined to all vertices in S , i.e. the intersection of the neighbourhoods of the vertices in S , and let $d(S) = |N(S)|$. For $S \in \mathcal{K}_k(G)$, let

$$f_G(S) = \sum_{T \subset S, |T|=k-1} d(T).$$

We begin by sketching the proof. The fact that the ratio between the K_k -density of G and the K_{k-1} -density of G is very close to $1/k$ will imply that the average $\mathbb{E}f_G(S)$ over all sets $S \in \mathcal{K}_k(G)$ is not too far below n . The fact that the K_{k+1} -density of G is small will mean that for most sets $S \in \mathcal{K}_k(G)$, every $(k-1)$ -subset $T \subset S$ has $N(T)$ spanning few edges of G , and any two distinct $(k-1)$ -subsets $T, T' \subset S$ have $|N(T) \cap N(T')|$ small. Hence, if we pick such a set S which has $f_G(S)$ not too far below the average, the sets $\{N(T) : T \subset S, |T| = k-1\}$ will be almost pairwise disjoint, will cover most of the vertices of G , and will each span few edges of G . Small alterations will produce a k -partition of $V(G)$ with few edges of G within each class, proving the theorem.

We now proceed with the proof. Observe that

$$\begin{aligned} \mathbb{E}f_G &= \frac{\sum_{S \in \mathcal{K}_k(G)} \sum_{T \subset S, |T|=k-1} d(T)}{K_k(G)} \\ &= \frac{\sum_{T \in \mathcal{K}_{k-1}(G)} d(T)^2}{K_k(G)} \\ &\geq \frac{\left(\sum_{T \in \mathcal{K}_{k-1}(G)} d(T)\right)^2}{K_{k-1}(G)K_k(G)} \\ &= \frac{(kK_k(G))^2}{K_{k-1}(G)K_k(G)} \\ &= k^2 \frac{K_k(G)}{K_{k-1}(G)} \\ &\geq k^2(1-\gamma) \frac{k!}{k^k} \frac{1}{1+\beta} \frac{k^{k-1}}{k!} \frac{\binom{n}{k}}{\binom{n}{k-1}} \\ &= \frac{1-\gamma}{1+\beta} (n-k+1). \end{aligned}$$

(The first inequality follows from Cauchy-Schwarz, and the second from our assumptions on the K_k -density and the K_{k-1} -density of G .)

We call a set $T \in \mathcal{K}_{k-1}(G)$ *dangerous* if it is contained in at least $\sqrt{\alpha} \binom{n-k+1}{2}$ K_{k+1} 's. Let D denote the number of dangerous $(k-1)$ -sets. Double-counting the number of times a $(k-1)$ -set is contained in a K_{k+1} , we obtain:

$$D\sqrt{\alpha} \binom{n-k+1}{2} \leq \binom{k+1}{2} \alpha \binom{n}{k+1},$$

since there are at most $\alpha \binom{n}{k+1}$ K_{k+1} 's in G . Hence,

$$D \leq \sqrt{\alpha} \binom{n}{k-1}.$$

Similarly, we call a set $S \in \mathcal{K}_k(G)$ *treacherous* if it is contained in at least $\sqrt{\alpha}(n-k)$ K_{k+1} 's. Double-counting the number of times a k -set is contained in a K_{k+1} , we see that there are at most $\sqrt{\alpha} \binom{n}{k}$ treacherous k -sets.

Call a set $S \in \mathcal{K}_k(G)$ *bad* if it is treacherous, or contains at least one dangerous $(k-1)$ -set; otherwise, call S *good*. Then the number of bad k -sets is at most

$$\sqrt{\alpha} \binom{n}{k} + (n-k+1)\sqrt{\alpha} \binom{n}{k-1} = (k+1)\sqrt{\alpha} \binom{n}{k},$$

so the fraction of sets in $\mathcal{K}_k(G)$ which are bad is at most

$$\frac{(k+1)\sqrt{\alpha}}{(1-\gamma)\frac{k!}{k^k}} = \frac{k^k(k+1)\sqrt{\alpha}}{(1-\gamma)k!}.$$

Suppose that

$$\max\{|f_G(S)| : S \text{ is good}\} < (1-\psi)(n-k+1).$$

Observe that for any $S \in \mathcal{K}_k(G)$, we have

$$f_G(S) \leq k(n-k+1),$$

since $d(T) \leq n-k+1$ for each $T \in S^{(k-1)}$. Hence,

$$\begin{aligned} \mathbb{E}f_G &< \left(\left(1 - \frac{k^k(k+1)\sqrt{\alpha}}{(1-\gamma)k!} \right) (1-\psi) + \frac{k^k(k+1)\sqrt{\alpha}}{(1-\gamma)k!} k \right) (n-k+1) \\ &\leq \left(1 - \psi + \frac{k^{k+1}(k+1)\sqrt{\alpha}}{(1-\gamma)k!} \right) (n-k+1), \end{aligned}$$

a contradiction if

$$\psi = \psi_0 := 1 - \frac{1-\gamma}{1+\beta} + \frac{k^{k+1}(k+1)\sqrt{\alpha}}{(1-\gamma)k!} \leq \gamma + \beta + \frac{2k^{k+1}(k+1)}{k!} \sqrt{\alpha}.$$

Let $S \in \mathcal{K}_k(G)$ be a good k -set such that $f_G(S) \geq (1 - \psi_0)(n - k + 1)$. Write $S = \{v_1, \dots, v_k\}$, let $T_i = S \setminus \{v_i\}$ for each i , and let $N_i = N(T_i)$ for each i . Observe that $N_i \cap N_j = N(S)$ for each $i \neq j$, and $|N(S)| = d(S) \leq \sqrt{\alpha}(n - k)$. Let $W_i = N_i \setminus N(S)$ for each i ; observe that the W_i 's are pairwise disjoint. Let

$$R = V(G) \setminus \cup_{i=1}^k W_i$$

be the set of ‘leftover’ vertices.

Observe that

$$\sum_{i=1}^k |N_i \setminus N(S)| = f_G(S) - kN(S) \geq (1 - \psi)(n - k + 1) - k\sqrt{\alpha}(n - k),$$

and therefore the number of leftover vertices satisfies

$$|R| < (\psi + k\sqrt{\alpha})n + k.$$

We now produce a k -partition $(V_i)_{i=1}^k$ of $V(G)$ by extending the partition $(W_i)_{i=1}^k$ of $V(G) \setminus R$ arbitrarily to R , i.e., we partition the leftover vertices arbitrarily. Now delete all edges of G within V_i for each i . The number of edges within N_i is precisely the number of K_{k+1} 's containing T_i , which is at most $\sqrt{\alpha} \binom{n-k+1}{2}$. The number of edges incident with R is trivially at most $(\psi + k\sqrt{\alpha})n(n - 1) + k(n - 1)$. Hence, the number of edges deleted was at most

$$\begin{aligned} & (\psi + k\sqrt{\alpha})n(n - 1) + k(n - 1) + k\sqrt{\alpha} \binom{n - k + 1}{2} \\ & \leq \left(2\beta + 2\gamma + \frac{8k^{k+1}(k + 1)}{k!} \sqrt{\alpha} + 2k/n \right) \binom{n}{2}. \end{aligned}$$

Removing an edge removes at most $\binom{n-2}{k-2}$ K_k 's, and therefore the total number of K_k 's removed is at most

$$\begin{aligned} & \left(2\beta + 2\gamma + \frac{8k^{k+1}(k + 1)}{k!} \sqrt{\alpha} + 2k/n \right) \binom{n}{2} \binom{n-2}{k-2} \\ & = \left(2\beta + 2\gamma + \frac{8k^{k+1}(k + 1)}{k!} \sqrt{\alpha} + 2k/n \right) \binom{k}{2} \binom{n}{k}, \end{aligned}$$

completing the proof. $\square$

Note that the two results above together imply the following

Corollary 15. *For any $k \in \mathbb{N}$, there exist constants $A_k, B_k > 0$ such that the following holds. For any $\alpha \geq 0$, if G is an n -vertex graph with K_{k+1} -density at most α , and K_k -density at least*

$$(1 - \gamma) \frac{k!}{k^k},$$

where $\gamma \leq 1/2$, then G can be made into a k -partite graph G_0 by removing at most

$$(2\gamma + A_k \alpha^{1/(k+2)} + B_k/n) \binom{n}{2}$$

edges, which removes at most

$$(2\gamma + A_k \alpha^{1/(k+2)} + B_k/n) \binom{k}{2} \binom{n}{k}$$

K_k 's.

Proof of Proposition 9. Suppose $\mathcal{G} \subset \mathcal{P}X$ has $|\mathcal{G}| = m \leq (1 + \eta)|\mathcal{F}_{n,k}|$, and k -generates at least $(1 - \epsilon)2^n$ subsets of X . Our aim is to show that $\mathcal{G}$ is close to a canonical k -generator. We may assume that $\epsilon \leq 1/C_k^k$ and $\eta \leq 1/D_k^k$, so by choosing C_k and D_k appropriately large, we may assume throughout that ϵ and η are small. By choosing ξ_k appropriately small, we may assume that $n \geq n_0(k)$, where $n_0(k)$ is any function of k .

We first apply Lemma 13 and Theorem 14 with $G = H[\mathcal{G}]$, where H is the Kneser graph on $\mathcal{P}X$, $\mathcal{G} \subset \mathcal{P}X$ with $|\mathcal{G}| = m \leq (1 + \eta)|\mathcal{F}_{n,k}|$, and $\mathcal{G}$ k -generates at least $(1 - \epsilon)2^n$ subsets of X . By (6), we have

$$d_{K_{k+1}}(H[\mathcal{G}]) \leq 2^{-a_k n},$$

and therefore we may take $\alpha = 2^{-a_k n}$. Applying Lemma 13 with $r = k - 1$, we may take $\beta = 2^{-b_k n}$ for some $b_k > 0$.

We have $|\mathcal{G}| = m \leq (1 + \eta)(k2^{n/k} - k)$, so

$$\binom{m}{k} \leq \frac{m^k}{k!} < \frac{(1 + \eta)^k k^k}{k!} 2^n.$$

Notice that

$$\sum_{i=0}^{k-1} \binom{m}{i} \leq km^{k-1} \leq k((1 + \eta)k2^{n/k})^{k-1} < (1 + \eta)^{k-1} k^k 2^{(1-1/k)n}.$$

Since $\mathcal{G}$ k -generates at least $(1 - \epsilon)2^n$ subsets of X , we have

$$K_k(H[\mathcal{G}]) \geq (1 - \epsilon)2^n - (1 + \eta)^{k-1} k^k 2^{(1-1/k)n}.$$

Hence,

$$\begin{aligned}
d_{K_k}(H[\mathcal{G}]) &= \frac{K_k(H[\mathcal{G}])}{\binom{m}{k}} \\
&\geq \frac{(1-\epsilon)2^n - (1+\eta)^{k-1}k^k 2^{(1-1/k)n}}{\binom{(1+\eta)^k 2^{n/k}}{k}} \\
&\geq \frac{1-\epsilon - (1+\eta)^{k-1}k^k 2^{-n/k}}{(1+\eta)^k} \frac{k!}{k^k} \\
&\geq (1-\epsilon - k\eta - k^k 2^{-n/k}) \frac{k!}{k^k},
\end{aligned}$$

where the last inequality follows from

$$\frac{1-\epsilon}{(1+\eta)^k} \geq (1-\epsilon)(1-\eta)^k \geq (1-\epsilon)(1-k\eta) \geq 1-\epsilon - k\eta.$$

Therefore, the K_k -density of $H[\mathcal{G}]$ satisfies

$$d_{K_k}(H[\mathcal{G}]) \geq (1-\gamma) \frac{k!}{k^k},$$

where

$$\gamma = \epsilon + k\eta + k^k 2^{-n/k}.$$

Let

$$\psi = \left(2\beta + 2\gamma + \frac{8k^{k+1}(k+1)}{k!} \sqrt{\alpha} + 2k/n \right) \binom{k}{2}.$$

By Theorem 14, there exists a k -partite subgraph G_0 of $H[\mathcal{G}]$ with

$$\begin{aligned}
K_k(G_0) &\geq K_k(H[\mathcal{G}]) - \psi \binom{m}{k} \\
&\geq (1-\epsilon)2^n - (1+\eta)^{k-1}k^k 2^{(1-1/k)n} - \psi \binom{m}{k} \\
&\geq \left(1-\epsilon - \frac{(1+\eta)^k k^k}{k!} \psi - (1+\eta)^{k-1}k^k 2^{-k/n} \right) 2^n.
\end{aligned}$$

Writing

$$\phi = \epsilon + \frac{(1+\eta)^k k^k}{k!} \psi + (1+\eta)^{k-1}k^k 2^{-k/n},$$

we have

$$K_k(G_0) \geq (1-\phi)2^n.$$

Let $V_1, \dots, V_k$ be the vertex-classes of G_0 . By the AM/GM inequality,

$$K_k(G_0) \leq \prod_{i=1}^k |V_i| \leq \left(\frac{\sum_{i=1}^k |V_i|}{k} \right)^k = (m/k)^k,$$

and therefore

$$|\mathcal{G}| = m \geq k(K_k(G_0))^{1/k} \geq k(1 - \phi)^{1/k} 2^{n/k}, \quad (8)$$

recovering the asymptotic result of [5].

Moreover, any k -partite graph G_0 satisfies

$$e(G_0) \geq \binom{k}{2} (K_k(G_0))^{2/k}.$$

To see this, simply apply Shearer's Entropy Lemma with $S = V(G_0)$, $\mathcal{F} = \mathcal{K}_k(G_0)$, and $\mathcal{A} = \{V_i \cup V_j : i \neq j\}$. Then $\mathcal{A}$ is a $(k-1)$ -cover of $V(G_0)$. Note that $\mathcal{F}_{V_i \cup V_j} \subset E_{G_0}(V_i, V_j)$, and therefore

$$(K_k(G_0))^{k-1} \leq \prod_{\{i,j\} \in [k]^{(2)}} e_{G_0}(V_i, V_j).$$

Applying the AM/GM inequality gives:

$$(K_k(G_0))^{k-1} \leq \prod_{\{i,j\}} e_{G_0}(V_i, V_j) \leq \left(\frac{\sum_{\{i,j\}} e_{G_0}(V_i, V_j)}{\binom{k}{2}} \right)^{\binom{k}{2}} = \left(\frac{e(G_0)}{\binom{k}{2}} \right)^{\binom{k}{2}},$$

and therefore

$$e(G_0) \geq \binom{k}{2} (K_k(G_0))^{2/k},$$

as required.

It follows that

$$\begin{aligned} e(G_0) &\geq \binom{k}{2} (1 - \phi)^{2/k} 2^{2n/k} \\ &\geq \binom{k}{2} (1 - \phi)^{2/k} \left(\frac{m}{(1 + \eta)k} \right)^2 \\ &\geq (1 - \eta)^2 (1 - \phi)^{2/k} (1 - 1/k) m^2 / 2 \\ &\geq (1 - 2\eta - \phi^{2/k}) (1 - 1/k) m^2 / 2 \\ &= (1 - \delta) (1 - 1/k) m^2 / 2, \end{aligned}$$

where $\delta = 2\eta + \phi^{2/k}$.

Hence, G_0 is a k -partite subgraph of $H[\mathcal{G}]$ with $|G_0| = |\mathcal{G}| = m$, and $e(G_0) \geq (1 - \delta - 1/k)m^2/2$. Applying Lemma 11 to G_0 , we see that there exists an induced subgraph H' of G_0 with

$$|H'| \geq (1 - \sqrt{\delta})|\mathcal{G}|, \quad (9)$$

and

$$\delta(H') \geq (1 - 1/k - \sqrt{\delta})(|H'| - 1).$$

Let $Y_1, \dots, Y_k$ be the vertex-classes of H' ; note that these are families of subsets of X . Clearly, for each $i \in [k]$,

$$|Y_i| \leq |H'| - \delta(H') \leq (1/k + \sqrt{\delta})|H'| + 1. \quad (10)$$

Hence, for each $i \in [k]$,

$$|Y_i| \geq |H'| - (k-1)((1/k + \sqrt{\delta})|H'| + 1) \geq (1/k - (k-1)\sqrt{\delta})|H'| - k + 1. \quad (11)$$

For each $i \in [k]$, let

$$S_i = \bigcup_{y \in Y_i} y$$

be the union of all sets in Y_i . We claim that the S_i 's are pairwise disjoint. Suppose for a contradiction that $S_1 \cap S_2 \neq \emptyset$. Then there exist $y_1 \in Y_1$ and $y_2 \in Y_2$ which both contain some element $p \in X$. Since

$$\delta(H') \geq (1 - 1/k - \sqrt{\delta})(|H'| - 1),$$

at least $(1 - 1/k - \sqrt{\delta})(|H'| - 1)$ sets in $\cup_{i \neq 1} Y_i$ do not contain p . By (10),

$$|\cup_{i \neq 1} Y_i| = \sum_{i \neq 1} |Y_i| \leq (1 - 1/k + (k-1)\sqrt{\delta})|H'| + k - 1,$$

and therefore the number of sets in $\cup_{i \neq 1} Y_i$ containing p is at most

$$(1 - 1/k + (k-1)\sqrt{\delta})|H'| + k - 1 - (1 - 1/k - \sqrt{\delta})(|H'| - 1) \leq k\sqrt{\delta}|H'| + k.$$

The same holds for the number of sets in $\cup_{i \neq 2} Y_i$ containing p , so the total number of sets in H' containing p is at most

$$2k\sqrt{\delta}|H'| + 2k.$$

Hence, the total number of sets in $\mathcal{G}$ containing p is at most

$$(2k + 1)\sqrt{\delta}m + 2k.$$

But then the number of ways of choosing at most k disjoint sets in $\mathcal{G}$ with one containing p is at most

$$(1 + m^{k-1})((2k+1)\sqrt{\delta}m + 2k) = O_k(\sqrt{\delta})2^n + O_k(2^{(1-1/k)n}) < 2^{n-1} - \epsilon 2^n,$$

contradicting the fact that $\mathcal{G}$ k -generates all but $\epsilon 2^n$ of the sets containing p .

Hence, we may conclude that the S_i 's are pairwise disjoint. By definition, $Y_i \subset \mathcal{P}S_i$, and therefore $|Y_i| \leq 2^{|S_i|}$. But from (11),

$$\begin{aligned} |Y_i| &\geq (1 - k(k-1)\sqrt{\delta})|H'|/k - k + 1 \\ &\geq (1 - k(k-1)\sqrt{\delta})(1 - \sqrt{\delta})|\mathcal{G}|/k - k + 1 \\ &\geq (1 - k(k-1)\sqrt{\delta})(1 - \sqrt{\delta})(1 - \phi)^{1/k}2^{n/k} - k + 1 \\ &\geq (1 - (k(k-1) + 1)\sqrt{\delta} - \phi^{1/k})2^{n/k} - k + 1 \\ &> (1 - k^2\sqrt{\delta} - \phi^{1/k})2^{n/k} - k \\ &> 2^{n/k-1}, \end{aligned}$$

using (9) and (8) for the second and third inequalities respectively. Hence, we must have $|S_i| \geq n/k$ for each i , and therefore $|S_i| = n/k$ for each i , i.e. $(S_i)_{i=1}^k$ is an equipartition of X . Putting everything together and recalling that $\delta = 2\eta + \phi^{2/k}$ and $\phi = O_k(\epsilon + \eta + 2^{-c_k n})$, we have

$$\begin{aligned} |\mathcal{G} \cap (\cup_{i=1}^k \mathcal{P}S_i)| &\geq \sum_{i=1}^k |Y_i| \\ &\geq (1 - k^2\sqrt{\delta} - \phi^{1/k})k2^{n/k} - k^2 \\ &\geq (1 - C_k\epsilon^{1/k} - D_k\eta^{1/k} - 2^{-\xi_k n})k2^{n/k} \end{aligned}$$

(provided n is sufficiently large depending on k), where $C_k, D_k, \xi_k > 0$ depend only on k . This proves Proposition 9. $\square$

We now prove the following

Proposition 16. *Let $\nu(n) = o(1)$. If $\mathcal{G}$ is a k -generator for X with $|\mathcal{G}| \leq |\mathcal{F}_{n,k}|$, and*

$$|\mathcal{G} \cap (\cup_{i=1}^k \mathcal{P}S_i)| \geq (1 - \nu)|\mathcal{F}_{n,k}|,$$

where $(S_i)_{i=1}^k$ is a partition of X into k classes of sizes as equal as possible, then provided n is sufficiently large depending on k , we have $|\mathcal{G}| = |\mathcal{F}_{n,k}|$ and

$$\mathcal{G} = \cup_{i=1}^k \mathcal{P}S_i \setminus \{\emptyset\}.$$

Note that n is no longer assumed to be a multiple of k ; the case $k = 2$ and n odd will be needed in Section 3.

Proof. Let $\mathcal{G}$ and $(S_i)_{i=1}^k$ be as in the statement of the proposition. For each $i \in [k]$, let $\mathcal{F}_i = (\mathcal{P}S_i \setminus \{\emptyset\}) \setminus \mathcal{G}$ be the collection of all nonempty subsets of S_i which are not in $\mathcal{G}$. By our assumption on $\mathcal{G}$, we know that $|\mathcal{F}_i| \leq o(2^{|S_i|})$ for each $i \in [k]$. Let

$$\mathcal{E} = \mathcal{G} \setminus \bigcup_{i=1}^k \mathcal{P}(S_i)$$

be the collection of ‘extra’ sets in $\mathcal{G}$; let $|\mathcal{E}| = M$.

By relabeling the S_i ’s, we may assume that $|\mathcal{F}_1| \geq |\mathcal{F}_2| \geq \dots \geq |\mathcal{F}_k|$. By our assumption on $|\mathcal{G}|$, $M \leq k|\mathcal{F}_1|$.

Let

$$\mathcal{R} = \{y_1 \sqcup s_2 \sqcup \dots \sqcup s_k : y_1 \in \mathcal{F}_1, s_i \subset S_i \ \forall i \geq 2\};$$

observe that the sets $y_1 \sqcup s_2 \sqcup \dots \sqcup s_k$ are all distinct, so $|\mathcal{R}| = |\mathcal{F}_1|2^{n-|S_1|}$. By considering the number of sets in $\mathcal{E}$ needed for $\mathcal{G}$ to k -generate $\mathcal{R}$, we will show that $M > k|\mathcal{F}_1|$ unless $\mathcal{F}_1 = \emptyset$. (In fact, our argument would also show that $M > p_k|\mathcal{F}_1|$ unless $\mathcal{F}_1 = \emptyset$, for any $p_k > 0$ depending only on k .)

Let N be the number of sets in $\mathcal{R}$ which may be expressed as a disjoint union of two sets in $\mathcal{E}$ and at most $k-2$ other sets in $\mathcal{G}$. Then

$$\begin{aligned} N &\leq \binom{M}{2} \sum_{i=0}^{k-2} \binom{m}{i} \\ &\leq \frac{1}{2}k^2|\mathcal{F}_1|^2(k-1) \frac{(c_0k2^{n/k})^{k-2}}{(k-2)!} \\ &\leq 4c_0^{k-2}k^k \left(\frac{|\mathcal{F}_1|}{2^{|S_1|}} \right) |\mathcal{F}_1|2^{n-|S_1|} \\ &= o(1)|\mathcal{F}_1|2^{n-|S_1|} \\ &= o(|\mathcal{R}|), \end{aligned} \tag{12}$$

where we have used $|\mathcal{G}| \leq |\mathcal{F}_{n,k}| \leq c_0k2^{n/k}$ (see (3)), $|S_1| \leq \lceil n/k \rceil$, and $|\mathcal{F}_1| = o(2^{|S_1|})$ in the second, third and fourth lines respectively.

Now fix $x_1 \in \mathcal{F}_1$. For $j \geq 1$, let $\mathcal{A}_j(x_1)$ be the collection of $(k-1)$ -tuples $(s_2, \dots, s_k) \in \mathcal{P}S_2 \times \dots \times \mathcal{P}S_k$ such that

$$x_1 \sqcup s_2 \sqcup \dots \sqcup s_k$$

may be expressed as a disjoint union

$$y_1 \sqcup y_2 \sqcup \dots \sqcup y_k$$

with $y_j \in \mathcal{E}$ but $y_i \subset S_i \ \forall i \neq j$. Let $\mathcal{A}^*(x_1)$ be the collection of $(k-1)$ -tuples $(s_2, \dots, s_k) \in \mathcal{P}S_2 \times \dots \times \mathcal{P}S_k$ such that

$$x_1 \sqcup s_2 \sqcup \dots \sqcup s_k$$

may be expressed as a disjoint union of two sets in $\mathcal{E}$ and at most $k-2$ other sets in $\mathcal{G}$.

Now fix $j \neq 1$. For each $(s_2, \dots, s_k) \in \mathcal{A}_j(x_1)$, we may write

$$x_1 \sqcup s_2 \sqcup \dots \sqcup s_k = s'_1 \sqcup s_2 \sqcup \dots \sqcup s_{j-1} \sqcup y_j \sqcup s_{j+1} \sqcup \dots \sqcup s_k,$$

where $y_j = s_j \sqcup (x_1 \setminus s'_1) \in \mathcal{E}$. Since $y_j \cap S_j = s_j$, different s_j 's correspond to different y_j 's $\in \mathcal{E}$, and so there are at most $|\mathcal{E}| = M$ choices for s_j . Therefore,

$$|\mathcal{A}_j(x_1)| \leq 2^{n-|S_1|-|S_j|} M \leq 2^{n-|S_1|-|S_j|} k |\mathcal{F}_1| \leq 2k \left(\frac{|\mathcal{F}_1|}{2^{|S_1|}} \right) 2^{n-|S_1|},$$

the last inequality following from the fact that $|S_j| \geq |S_1| - 1$. Hence,

$$\sum_{j=2}^k |\mathcal{A}_j(x_1)| \leq 2k(k-1) \left(\frac{|\mathcal{F}_1|}{2^{|S_1|}} \right) 2^{n-|S_1|} = o(1) 2^{n-|S_1|}. \quad (13)$$

Observe that for each $x_1 \in \mathcal{F}_1$,

$$\mathcal{A}^*(x_1) \cup \bigcup_{j=1}^k \mathcal{A}_j(x_1) = \mathcal{P}S_2 \times \mathcal{P}S_3 \times \dots \times \mathcal{P}S_k,$$

and therefore

$$|\mathcal{A}^*(x_1)| + |\mathcal{A}_1(x_1)| + \sum_{j=2}^k |\mathcal{A}_j(x_1)| \geq 2^{n-|S_1|},$$

so by (13),

$$|\mathcal{A}^*(x_1)| + |\mathcal{A}_1(x_1)| \geq (1 - o(1)) 2^{n-|S_1|}.$$

Call $x_1 \in \mathcal{F}_1$ ‘bad’ if $|\mathcal{A}^*(x_1)| \geq 2^{-(k+2)} 2^{n-|S_1|}$; otherwise, call x_1 ‘good’. By (12), at most a $o(1)$ -fraction of the sets in $\mathcal{F}_1$ are bad, so at least a $1 - o(1)$ fraction are good. For each good set $x_1 \in \mathcal{F}_1$, notice that

$$|\mathcal{A}_1(x_1)| \geq (1 - 2^{-(k+2)} - o(1)) 2^{n-|S_1|}.$$

Now perform the following process. Choose any $(s_2, \dots, s_k) \in \mathcal{A}_1(x_1)$; we may write

$$x_1 \sqcup s_2 \sqcup \dots \sqcup s_k = z^{(1)} \sqcup s'_2 \sqcup \dots \sqcup s'_k$$

with $(s'_2, \dots, s'_k) \in \mathcal{P}S_2 \times \dots \times \mathcal{P}S_k$, $z^{(1)} \in \mathcal{E}$, $z^{(1)} \cap S_1 = x_1$, and $z^{(1)} \setminus S_1 \neq \emptyset$. Pick $p_1 \in z^{(1)} \setminus S_1$. At most $\frac{1}{2} 2^{n-|S_1|}$ of the members of $\mathcal{A}_1(x_1)$ have union containing p_1 , so there are at least

$$(1 - \frac{1}{2} - 2^{-(k+2)} - o(1)) 2^{n-|S_1|}$$

remaining members of $\mathcal{A}_1(x_1)$. Choose one of these, $(t_2, \dots, t_k)$ say. By definition, we may write

$$x_1 \sqcup t_2 \sqcup \dots \sqcup t_k = z^{(2)} \sqcup t'_2 \sqcup \dots \sqcup t'_k$$

with $(t'_2, \dots, t'_k) \in \mathcal{P}S_2 \times \dots \times \mathcal{P}S_k$, $z^{(2)} \in \mathcal{E}$, $z^{(2)} \cap S_1 = x_1$, and $z^{(2)} \setminus S_1 \neq \emptyset$. Since $p_1 \notin z^{(2)}$, we must have $z^{(2)} \neq z^{(1)}$. Pick $p_2 \in z^{(2)} \setminus S_1$, and repeat. At most $\frac{3}{4}2^{n-|S_1|}$ of the members of $\mathcal{A}_1(x_1)$ have union containing p_1 or p_2 ; there are at least

$$(\frac{1}{4} - 2^{-(k+2)} - o(1))2^{n-|S_1|}$$

members remaining. Choose one of these, $(u_2, \dots, u_k)$ say. By definition, we may write

$$x_1 \sqcup u_2 \sqcup \dots \sqcup u_k = z^{(3)} \sqcup u'_2 \sqcup \dots \sqcup u'_k$$

with $(u'_2, \dots, u'_k) \in \mathcal{P}S_2 \times \dots \times \mathcal{P}S_k$, $z^{(3)} \in \mathcal{E}$, $z^{(3)} \cap S_1 = x_1$, and $z^{(3)} \setminus S_1 \neq \emptyset$. Note that again $z^{(3)}$ is distinct from $z^{(1)}, z^{(2)}$, since $p_1, p_2 \notin z^{(3)}$. Continuing this process for $k+1$ steps, we end up with a collection of $k+1$ distinct sets $z^{(1)}, \dots, z^{(k+1)} \in \mathcal{E}$ such that $z^{(l)} \cap S_1 = x_1 \forall l \in [k+1]$. Do this for each good set $x_1 \in \mathcal{F}_1$; the collections produced are clearly pairwise disjoint. Therefore,

$$|\mathcal{E}| \geq (k+1)(1 - o(1))|\mathcal{F}_1|.$$

This is a contradiction, unless $\mathcal{F}_1 = \emptyset$. Hence, we must have $\mathcal{F}_2 = \dots = \mathcal{F}_k = \emptyset$, and therefore

$$\mathcal{G} = \cup_{i=1}^k \mathcal{P}(S_i) \setminus \{\emptyset\},$$

proving Proposition 16, and completing the proof of Theorem 5. $\square$

3 The case $k = 2$ via bipartite subgraphs of H .

Our aim in this section is to prove the $k = 2$ case of Conjecture 1 for all sufficiently large *odd* n , which together with the $k = 2$ case of Theorem 5 will imply

Theorem 4. *If n is sufficiently large, X is an n -element set, and $\mathcal{G} \subset \mathcal{P}X$ is a 2-generator for X , then $|\mathcal{G}| \geq |\mathcal{F}_{n,2}|$. Equality holds only if $\mathcal{G}$ is of the form $\mathcal{F}_{n,2}$.*

Recall that

$$|\mathcal{F}_{n,2}| = \begin{cases} 2 \cdot 2^{n/2} - 2 & \text{if } n \text{ is even;} \\ 3 \cdot 2^{(n-1)/2} - 2 & \text{if } n \text{ is odd.} \end{cases}$$

Suppose that X is an n -element set, and $\mathcal{G} \subset \mathcal{P}X$ is a 2-generator for X with $|\mathcal{G}| = m \leq |\mathcal{F}_{n,2}|$. The counting argument in the Introduction gives

$$1 + m + \binom{m}{2} \geq 2^n,$$

which implies that

$$|\mathcal{G}| \geq (1 - o(1))\sqrt{2}2^{n/2}.$$

For n odd, we wish to improve this bound by a factor of approximately 1.5.

Our first aim is to prove that induced subgraphs of the Kneser graph H which have order $\Omega(2^{n/2})$ are $o(1)$ -close to being bipartite (Proposition 18).

Recall that a graph $G = (V, E)$ is said to be ϵ -close to being bipartite if it can be made bipartite by the removal of at most $\epsilon|V|^2$ edges, and ϵ -far from being bipartite if it requires the removal of at least $\epsilon|V|^2$ edges to make it bipartite.

Using Szemerédi's Regularity Lemma, Bollobás, Erdős, Simonovits and Szemerédi [3] proved the following.

Theorem 17 (Bollobás, Erdős, Simonovits, Szemerédi). *For any $\epsilon > 0$, there exists $g(\epsilon) \in \mathbb{N}$ depending on ϵ alone such that for any graph G which is ϵ -far from being bipartite, the probability that a uniform random induced subgraph of G of order $g(\epsilon)$ is non-bipartite is at least $1/2$.*

Building on methods of Goldreich, Goldwasser and Ron [12], Alon and Krivelevich [2] proved without using the Regularity Lemma that in fact, one may take

$$g(\epsilon) \leq \frac{(\log(1/\epsilon))^b}{\epsilon} \tag{14}$$

where $b > 0$ is an absolute constant. As observed in [2], this is tight up to the poly-logarithmic factor, since necessarily,

$$g(\epsilon) \geq \frac{1}{6\epsilon}.$$

We will first show that for any fixed $c > 0$ and $l \in \mathbb{N}$, if $\mathcal{A} \subset \mathcal{P}X$ with $|\mathcal{A}| \geq c2^{n/2}$, then the density of C_{2l+1} 's in $H[\mathcal{A}]$ is at most $o(1)$. To prove this, we will show that for any $l \in \mathbb{N}$, there exists $t \in \mathbb{N}$ such that for any fixed $c > 0$, if $\mathcal{A} \subset \mathcal{P}X$ with $|\mathcal{A}| \geq c2^{n/2}$, then the homomorphism density of $C_{2l+1} \otimes t$ in $H[\mathcal{A}]$ is $o(1)$. Using Lemma 7, we will deduce that the homomorphism density of C_{2l+1} in $H[\mathcal{A}]$ is $o(1)$, implying that the density of C_{2l+1} 's in $H[\mathcal{A}]$ is $o(1)$. This will show that $H[\mathcal{A}]$ is $o(1)$ -close to being bipartite (Proposition 18). To obtain a sharper estimate for the $o(1)$ term

in Proposition 18, we will use (14), although to prove Theorem 4, any $o(1)$ term would suffice, so one could in fact use Theorem 17 instead of (14).

We are now ready to prove the following

Proposition 18. *Let $c > 0$. Then there exists $b > 0$ such that for any $\mathcal{A} \subset \mathcal{P}X$ with $|\mathcal{A}| \geq c2^{n/2}$, the induced subgraph $H[\mathcal{A}]$ can be made bipartite by removing at most*

$$\frac{(\log_2 \log_2 n)^b}{\log_2 n} |\mathcal{A}|^2$$

edges.

Proof. Fix $c > 0$; let $\mathcal{A} \subset \mathcal{P}X$ with $|\mathcal{A}| = m \geq c2^{n/2}$. First, we show that for any fixed $l \in \mathbb{N}$, there exists $t \in \mathbb{N}$ such that the homomorphism density of $C_{2l+1} \otimes t$'s in $H[\mathcal{A}]$ is at most $o(1)$. The argument is a strengthening of that used by Alon and Frankl to prove Lemma 4.2 in [1].

Let $t \in \mathbb{N}$ to be chosen later. Choose $(2l+1)t$ members of $\mathcal{A}$ uniformly at random with replacement, $(A_i^{(j)})_{1 \leq i \leq 2l+1, 1 \leq j \leq t}$. The homomorphism density of $C_{2l+1} \otimes t$ in $H[\mathcal{A}]$ is precisely the probability that the unions

$$U_i = \bigcup_{j=1}^t A_i^{(j)}$$

satisfy $U_i \cap U_{i+1} = \emptyset$ for each i (where the addition is modulo $2l+1$).

We claim that if this occurs, then $|U_i| < (\frac{1}{2} - \eta)n$ for some i , provided $\eta < 1/(4l+2)$. Suppose for a contradiction that $U_i \cap U_{i+1} = \emptyset$ for each i , and $|U_i| \geq (\frac{1}{2} - \eta)n$ for each i . Then $|U_{i+2} \setminus U_i| \leq n - |U_{i+1}| - |U_i| \leq 2\eta n$ for each $i \in [2l-1]$. Since $U_{2l+1} \setminus U_1 \subset \bigcup_{j=1}^l (U_{2j+1} \setminus U_{2j-1})$, we have $|U_{2l+1} \setminus U_1| \leq \sum_{j=1}^l |U_{2j+1} \setminus U_{2j-1}| \leq 2l\eta n$. It follows that $|U_1 \cap U_{2l+1}| \geq (1/2 - (2l+1)\eta)n > 0$ if $\eta < 1/(4l+2)$, a contradiction.

We now show that the probability of this event is very small. Fix $i \in [k]$.

Observe that

$$\begin{aligned}
\Pr\{|U_i| \leq (1/2 - \eta)n\} &= \Pr\left(\bigcup_{S \subset X: |S| \leq (1/2 - \eta)n} \left(\bigcap_{j=1}^t \{A_i^{(j)} \subset S\}\right)\right) \\
&\leq \sum_{|S| \leq (1/2 - \eta)n} \Pr\left(\bigcap_{j=1}^t \{A_i^{(j)} \subset S\}\right) \\
&= \sum_{|S| \leq (1/2 - \eta)n} (2^{|S|}/m)^t \\
&\leq 2^n \left(\frac{2^{(1/2 - \eta)n}}{c^{2n/2}}\right)^t \\
&= 2^{-(\eta t - 1)n} c^{-t} \\
&\leq 2^{-n} c^{-t},
\end{aligned}$$

provided $t \geq 2/\eta$. Hence,

$$\Pr\left(\bigcup_{i=1}^{2l+1} \{|U_i| \leq (1/2 - \eta)n\}\right) \leq \sum_{i=1}^{2l+1} \Pr\{|U_i| \leq (1/2 - \eta)n\} \leq (2l+1)2^{-n} c^{-t}.$$

Therefore,

$$h_{C_{2l+1} \otimes t}(H[\mathcal{A}]) \leq (2l+1)2^{-n} c^{-t}.$$

Choose $\eta = \frac{1}{8l}$ and $t = 2/\eta = 16l$. By Lemma 7,

$$\begin{aligned}
h_{C_{2l+1}}(H[\mathcal{A}]) &\leq ((2l+1)2^{-n} c^{-t})^{1/t^{2l+1}} \\
&= (2l+1)^{1/(16l)^{2l+1}} 2^{-n/(16l)^{2l+1}} c^{-1/(16l)^{2l}} \\
&= O(2^{-n/(16l)^{2l+1}}).
\end{aligned}$$

Observe that the number of $(2s+1)$ -subsets of $\mathcal{A}$ containing an odd cycle of H is at most

$$\sum_{l=1}^s m^{2l+1} h_{C_{2l+1}}(H[\mathcal{A}]) \binom{m - (2l+1)}{2(s-l)}.$$

Hence, the probability that a uniform random $(2s+1)$ -subset of $\mathcal{A}$ contains an odd cycle of H is at most

$$\begin{aligned}
\sum_{l=1}^s \frac{m^{2l+1}}{m(m-1) \cdots (m-2l)} (2s+1)(2s) \cdots (2(s-l)+1) h_{C_{2l+1}}(H[\mathcal{A}]) \\
\leq s(2s+1)! O(2^{-n/(16s)^{2s+1}}),
\end{aligned}$$

(provided $s \leq O(\sqrt{m})$). This can be made $< 1/2$ by choosing

$$s = a \log_2 n / \log_2 \log_2 n,$$

for some suitable $a > 0$ depending only on c . By (14), it follows that $H[\mathcal{A}]$ is $((\log_2 \log_2 n)^b / \log_2 n)$ -close to being bipartite, for some suitable $b > 0$ depending only on c , proving the proposition. $\square$

Before proving Theorem 4 for n odd, we need some more definitions. Let X be a finite set. If $A \subset \mathcal{P}X$, and $i \in X$, we define

$$\begin{aligned} A_i^- &= \{x \in A : i \notin x\}, \\ A_i^+ &= \{x \setminus \{i\} : x \in A, i \in x\}; \end{aligned}$$

these are respectively called the *lower* and *upper i -sections* of A .

If Y and Z are disjoint subsets of X , we write $H[Y, Z]$ for the bipartite subgraph of the Kneser graph H consisting of all edges between Y and Z . If B is a bipartite subgraph of H with vertex-sets Y and Z , and $\mathcal{F} \subset \mathcal{P}X$, we say that B *2-generates* $\mathcal{F}$ if for every set $x \in \mathcal{F}$, there exist $y \in Y$ and $z \in Z$ such that $y \cap z = \emptyset$, $yz \in E(B)$, and $y \sqcup z = x$, i.e. every set in $\mathcal{F}$ corresponds to an edge of B .

Proof of Theorem 4 for n odd. Suppose that $n = 2l + 1 \geq 3$ is odd, X is an n -element set, and $\mathcal{G} \subset \mathcal{P}X$ is a 2-generator for X with $|\mathcal{G}| = m \leq |\mathcal{F}_{n,2}| = 3 \cdot 2^l - 2$. Observe that

$$e(H[\mathcal{G}]) \geq 2^{2l+1} - |\mathcal{G}| - 1 \geq 2^{2l+1} - 3 \cdot 2^l + 1,$$

and therefore $H[\mathcal{G}]$ has edge-density at least

$$\frac{2^{2l+1} - 3 \cdot 2^l + 1}{\binom{|\mathcal{G}|}{2}} \geq \frac{2^{2l+1} - 3 \cdot 2^l + 1}{\frac{1}{2}(3 \cdot 2^l - 2)(3 \cdot 2^l - 3)} > \frac{4}{9}.$$

(Here, the last inequality rearranges to the statement $l > 0$.) By Proposition 18 applied to $\mathcal{G}$, we can remove at most

$$\frac{(\log_2 \log_2 n)^b}{\log_2 n} |\mathcal{G}|^2 < \frac{(\log_2 \log_2 n)^b}{\log_2 n} 9 \cdot 2^{2l}$$

edges from $H[\mathcal{G}]$ to produce a bipartite graph B . Let Y, Z be the vertex-classes of B ; we may assume that $Y \sqcup Z = \mathcal{G}$. Define $\epsilon > 0$ by

$$|\{y \sqcup z : y \in Y, z \in Z, y \cap z = \emptyset\}| = (1 - \epsilon) 2^{2l+1};$$

then clearly, we have

$$e(B) \geq (1 - \epsilon)2^{2l+1}. \quad (15)$$

Note that

$$\epsilon \leq \frac{9}{2} \frac{(\log_2 \log_2 n)^b}{\log_2 n} + 3 \cdot 2^{-(l+1)} = O\left(\frac{(\log_2 \log_2 n)^b}{\log_2 n}\right) = o(1).$$

Let

$$\alpha = |Y|/2^l, \beta = |Z|/2^l.$$

By assumption, $\alpha + \beta \leq 3 - 2^{-(l-1)} < 3$. Since $|Y||Z| \geq e(B) \geq (2 - 2\epsilon)2^{2l}$, we have $\alpha\beta \geq 2 - 2\epsilon$. This implies that

$$1 - 2\epsilon < \alpha, \beta < 2 + 2\epsilon. \quad (16)$$

(To see this, simply observe that to maximize $\alpha\beta$ subject to the conditions $\alpha \leq 1 - 2\epsilon$ and $\alpha + \beta \leq 3$, it is best to take $\alpha = 1 - 2\epsilon$ and $\beta = 2 + 2\epsilon$, giving $\alpha\beta = 2 - 2\epsilon - 4\epsilon^2 < 2 - 2\epsilon$, a contradiction. It follows that we must have $\alpha > 1 - 2\epsilon$, so $\beta < 2 + 2\epsilon$; (16) follows by symmetry.)

From now on, we think of X as the set $[n] = \{1, 2, \dots, n\}$. Let

$$\begin{aligned} W_1 &= \{i \in [n] : |Y_i^+| \geq |Y|/3\}, \\ W_2 &= \{i \in [n] : |Z_i^+| \geq |Z|/3\}. \end{aligned}$$

First, we prove the following

Claim 1. $W_1 \cup W_2 = [n]$.

Proof. Suppose for a contradiction that $W_1 \cup W_2 \neq [n]$. Without loss of generality, we may assume that $n \notin W_1 \cup W_2$. Let

$$\theta = |Y_n^+|/|Y|, \phi = |Z_n^+|/|Z|;$$

then we have $\theta, \phi \leq 1/3$. Observe that the number e_n of edges between Y and Z which generate a set containing n satisfies

$$(1 - 2\epsilon)2^{2l} \leq e_n \leq (\theta\alpha(1 - \phi)\beta + \phi\beta(1 - \theta)\alpha)2^{2l} = (\theta + \phi - 2\theta\phi)\alpha\beta 2^{2l}. \quad (17)$$

(Here, the left-hand inequality comes from the fact that B 2-generates all but at most $\epsilon 2^{2l+1}$ subsets of $[n]$, and therefore B 2-generates at least $(1 - 2\epsilon)2^{2l}$ sets containing n .)

Notice that the function

$$f(\theta, \phi) = \theta + \phi - 2\theta\phi, \quad 0 \leq \theta, \phi \leq 1/3$$

is a strictly increasing function of both θ and ϕ for $0 \leq \theta, \phi \leq 1/3$, and therefore attains its maximum of $4/9$ at $\theta = \phi = 1/3$. Therefore,

$$1 - 2\epsilon \leq \frac{4}{9}\alpha\beta;$$

since $\alpha + \beta \leq 3$, we have

$$3/2 - 3\sqrt{\epsilon/2} \leq \alpha, \beta \leq 3/2 + 3\sqrt{\epsilon/2}.$$

Moreover, by the AM/GM inequality, $\alpha\beta \leq 9/4$, so

$$1 - 2\epsilon \leq \frac{9}{4}f(\theta, \phi), \quad (18)$$

and therefore

$$1/3 - 8\epsilon/3 \leq \theta, \phi \leq 1/3.$$

Thus $|Y|, |Z| = (3/2 - o(1))2^l$ and $\theta, \phi = 1/3 - o(1)$. Therefore, we have

$$\begin{aligned} |Y_n^+| &= 2^{l-1}(1 - o(1)), \\ |Z_n^+| &= 2^{l-1}(1 - o(1)), \\ |Y_n^-| &= 2^l(1 + o(1)), \\ |Z_n^-| &= 2^l(1 + o(1)). \end{aligned}$$

Observe that $\mathcal{G}_n^- = Y_n^- \cup Z_n^-$ must 2-generate all but at most $o(2^{2l})$ of the sets in $\mathcal{P}\{1, 2, \dots, n-1\} = \mathcal{P}\{1, 2, \dots, 2l\}$, and therefore, by Proposition 9 for $k = 2$ and n even, there exists an equipartition $S_1 \cup S_2$ of $\{1, 2, \dots, 2l\}$ such that Y_n^- contains at least $(1 - o(1))2^l$ members of $\mathcal{P}S_1$, and Z_n^- contains at least $(1 - o(1))2^l$ members of $\mathcal{P}S_2$. Define

$$\begin{aligned} U &= \{y \in Y : y \cap S_2 = \emptyset\}, \\ V &= \{z \in Z : z \cap S_1 = \emptyset\}. \end{aligned}$$

Since $|U_n^-| = (1 - o(1))2^l$ and $|V_n^-| = (1 - o(1))2^l$, we must have $|Y_n^- \setminus U_n^-| = o(2^l)$, and $|Z_n^- \setminus V_n^-| = o(2^l)$. Our aim is now to show that $|Y_n^+ \setminus U_n^+| = o(2^l)$, and $|Z_n^+ \setminus V_n^+| = o(2^l)$.

Clearly, we have $U_n^- \subset \mathcal{P}S_1$, and $V_n^- \subset \mathcal{P}S_2$, so $|U_n^-| \leq 2^l$ and $|V_n^-| \leq 2^l$. Moreover, each set $x \in Y_n^+ \setminus U_n^+$ contains an element of S_2 , and therefore $x \cup \{n\}$ is disjoint from at most 2^{l-1} sets in $V_n^- \subset \mathcal{P}S_2$. Similarly, each set $x \in Z_n^+ \setminus V_n^+$ contains an element of S_1 , and therefore $x \cup \{n\}$ is disjoint from at most 2^{l-1} sets in $U_n^- \subset \mathcal{P}S_1$. It follows that

$$\begin{aligned} e_n &\leq |U_n^+||V_n^-| + |Y_n^+ \setminus U_n^+|2^{l-1} + |V_n^+||U_n^-| + |Z_n^+ \setminus V_n^+|2^{l-1} \\ &\quad + |Y_n^- \setminus U_n^-||Z_n^+| + |Z_n^- \setminus V_n^-||Y_n^+| \\ &\leq |U_n^+|2^l + |Y_n^+ \setminus U_n^+|2^{l-1} + |V_n^+|2^l + |Z_n^+ \setminus V_n^+|2^{l-1} + o(2^{2l}). \end{aligned}$$

On the other hand, by (17), we have $e_n \geq (1 - o(1))2^{2l}$. Since $|Y_n^+| = 2^{l-1}(1 - o(1))$, and $|Z_n^+| = 2^{l-1}(1 - o(1))$, we must have $|Y_n^+ \setminus U_n^+| = o(2^l)$, and $|Z_n^+ \setminus V_n^+| = o(2^l)$, as required.

We may conclude that $|Y \setminus U| = o(2^l)$ and $|Z \setminus V| = o(2^l)$. Hence, there are at most $o(2^l)$ sets in $Y \cup Z = \mathcal{G}$ that intersect both S_1 and S_2 . On the other hand, since $|Y_n^+| = (1 - o(1))2^{l-1}$ and $|Z_n^+| = (1 - o(1))2^{l-1}$, there are at least $(1 + o(1))2^{l-1}$ sets $s_1 \subset S_1$ such that $s_1 \cup \{n\} \notin Y$, and there are at least $(1 + o(1))2^{l-1}$ sets $s_2 \subset S_2$ such that $s_2 \cup \{n\} \notin Z$. Taking all pairs s_1, s_2 gives at least $(1 + o(1))2^{2l-2}$ sets of the form

$$\{n\} \cup s_1 \cup s_2 \quad (s_1 \subset S_1, s_1 \cup \{n\} \notin Y, s_2 \subset S_2, s_2 \cup \{n\} \notin Z). \quad (19)$$

Each of these requires a set intersecting both S_1 and S_2 to express it as a disjoint union of two sets from $\mathcal{G}$. Since there are $o(2^l)$ members of $\mathcal{G}$ intersecting both S_1 and S_2 , $\mathcal{G}$ generates at most

$$(|\mathcal{G}| + 1)o(2^l) = o(2^{2l})$$

sets of the form (19), a contradiction. This proves the claim. $\square$

We now prove the following

Claim 2. $W_1 \cap W_2 = \emptyset$.

Proof. Suppose for a contradiction that $W_1 \cap W_2 \neq \emptyset$. Without loss of generality, we may assume that $n \in W_1 \cap W_2$. As before, let

$$\theta = |Y_n^+|/|Y|, \quad \phi = |Z_n^+|/|Z|;$$

this time, we have $\theta, \phi \geq 1/3$. Observe that

$$(2 - 2\epsilon)2^{2l} \leq e(B) \leq (1 - \theta\phi)\alpha\beta 2^{2l}. \quad (20)$$

Here, the left-hand inequality is (15), and the right-hand inequality comes from the fact that there are no edges between pairs of sets $(y, z) \in Y \times Z$ such that $n \in y \cap z$. Since $1 - \theta\phi \leq 8/9$, we have

$$2 - 2\epsilon \leq \frac{8}{9}\alpha\beta.$$

Since $\alpha + \beta \leq 3$, it follows that

$$\frac{3}{2}(1 - \sqrt{\epsilon}) \leq \alpha, \beta \leq \frac{3}{2}(1 + \sqrt{\epsilon}).$$

Since $\alpha\beta \leq 9/4$, we have

$$2 - 2\epsilon \leq \frac{9}{4}(1 - \theta\phi),$$

and therefore

$$1/3 \leq \theta, \phi \leq 1/3 + 8\epsilon/3.$$

Hence, we have

$$\begin{aligned} |Y_n^+| &= 2^{l-1}(1 - o(1)), \\ |Z_n^+| &= 2^{l-1}(1 - o(1)), \\ |Y_n^-| &= 2^l(1 + o(1)), \\ |Z_n^-| &= 2^l(1 + o(1)), \end{aligned}$$

so exactly as in the proof of Claim 1, we obtain a contradiction. $\square$

Claims 1 and 2 together imply that $W_1 \cup W_2$ is a partition of $\{1, 2, \dots, n\} = \{1, 2, \dots, 2l + 1\}$. We will now show that at least a $(2/3 - o(1))$ -fraction of the sets in Y are subsets of W_1 , and similarly at least a $(2/3 - o(1))$ -fraction of the sets in Z are subsets of W_2 . Let

$$\sigma = \frac{|Y \setminus \mathcal{P}(W_1)|}{|Y|}, \quad \tau = \frac{|Z \setminus \mathcal{P}(W_2)|}{|Z|}.$$

Let $y \in Y \setminus \mathcal{P}(W_1)$, and choose $i \in y \cap W_2$; since at least $|Z|/3$ of the sets in Z contain i , y has at most $2|Z|/3$ neighbours in Z . Hence,

$$(2 - 2\epsilon)2^{2l} \leq e(B) \leq (\tfrac{2}{3}\sigma\alpha\beta + (1 - \sigma)\alpha\beta)2^{2l} = (1 - \sigma/3)\alpha\beta 2^{2l} \leq (1 - \sigma/3)\tfrac{9}{4}2^{2l}, \quad (21)$$

and therefore

$$\sigma \leq 1/3 + 8\epsilon/3,$$

so

$$|Y \cap \mathcal{P}(W_1)| \geq (2/3 - 8\epsilon/3)|Y|. \quad (22)$$

Similarly, $\tau \leq 1/3 + 8\epsilon/3$, and therefore $|Z \cap \mathcal{P}(W_2)| \geq (2/3 - 8\epsilon/3)|Z|$.

If $|W_1| \leq l - 1$, then $|Y \cap \mathcal{P}(W_1)| \leq 2^{l-1}$, so

$$|Y| \leq \frac{2^{l-1}}{2/3 - 8\epsilon/3} = \tfrac{3}{4} \frac{2^l}{1 - 4\epsilon} < (1 - 2\epsilon)2^l,$$

contradicting (16). Hence, we must have $|W_1| \geq l$. Similarly, $|W_2| \geq l$, so $\{|W_1|, |W_2|\} = \{l, l + 1\}$. Without loss of generality, we may assume that $|W_1| = l$ and $|W_2| = l + 1$.

We now observe that

$$|Z| \geq (3/2 - 6\epsilon)2^l \quad (23)$$

To see this, suppose that $|Z| = (3/2 - \eta)2^l$. Since $|Z| + |Y| < 3 \cdot 2^l$, we have $|Y| \leq (3/2 + \eta)2^l$. Recall that any $y \in Y \setminus \mathcal{P}W_1$ has at most $2|Z|/3$ neighbours in Z . Thus, we have

$$\begin{aligned} (2 - 2\epsilon)2^{2l} &\leq e(B) \\ &\leq |Y \cap \mathcal{P}W_1||Z| + |Y \setminus \mathcal{P}W_1|\frac{2}{3}|Z| \\ &\leq 2^l(\frac{3}{2} - \eta)2^l + (\frac{1}{2} + \eta)2^l\frac{2}{3}(\frac{3}{2} - \eta)2^l \\ &= (2 - \frac{1}{3}\eta - \frac{2}{3}\eta^2)2^{2l}. \end{aligned}$$

Therefore $\eta \leq 6\epsilon$, i.e. $|Z| \geq (3/2 - 6\epsilon)2^l$, as claimed. Since $|Z| + |Y| < 3 \cdot 2^l$, we have

$$|Y| \leq (3/2 + 6\epsilon)2^l. \quad (24)$$

We now prove the following

Claim 3. (a) $|\mathcal{P}(W_1) \setminus Y| \leq 22\epsilon 2^l$;
(b) $|Z \setminus \mathcal{P}W_2| \leq (\sqrt{\epsilon} + 2\epsilon)2^l$.

Proof. We prove this by constructing another bipartite subgraph B_2 of H with the same number of vertices as B , and comparing $e(B_2)$ with $e(B)$. First, let

$$D = \min\{|\mathcal{P}(W_2) \setminus Z|, |Z \setminus \mathcal{P}W_2|\},$$

add D new members of $\mathcal{P}(W_2) \setminus Z$ to Z , and delete D members of $Z \setminus \mathcal{P}W_2$, producing a new set Z' and a new bipartite graph $B_1 = H[Y, Z']$. Since $|Z'| = |Z| \leq (2 + 2\epsilon)2^l$, we have $|Z' \setminus \mathcal{P}W_2| \leq \epsilon 2^{l+1}$, i.e. Z' is almost contained within $\mathcal{P}W_2$. Notice that every member $z \in Z \setminus \mathcal{P}W_2$ had at most $2|Y|/3$ neighbours in Y , and every new member of Z' has at least $|Y \cap \mathcal{P}(W_1)| \geq (2/3 - 8\epsilon/3)|Y|$ neighbours in Y , using (22). Hence,

$$e(B) - e(B_1) \leq \frac{8\epsilon}{3}|Y|D \leq \frac{8\epsilon}{3}|Y|\frac{2}{3}|Z| \leq \frac{16\epsilon}{9}\frac{9}{4}2^{2l} = 4\epsilon 2^{2l},$$

and therefore

$$e(B_1) \geq e(B) - 2\epsilon 2^{2l+1} \geq (1 - 3\epsilon)2^{2l+1}.$$

Second, let

$$C = \min\{|\mathcal{P}W_1 \setminus Y|, |Y \setminus \mathcal{P}W_1|\},$$

add C new members of $\mathcal{P}(W_1) \setminus Y$ to Y , and delete C members of $Y \setminus \mathcal{P}W_1$, producing a new set Y' and a new bipartite graph $B_2 = H[Y', Z']$. Since $|Y| \geq (1 - 2\epsilon)2^l$, we have $|Y' \cap \mathcal{P}W_1| \geq (1 - 2\epsilon)2^l$. Since every deleted member of Y contained an element of W_2 , it had at most $(1 + 2\epsilon)2^l$ neighbours in Z' . (Indeed, such member of Y intersects 2^l sets in $\mathcal{P}W_2$, so has at most 2^l neighbours in $Z' \cap \mathcal{P}W_2$; there are $|Z' \setminus \mathcal{P}W_2| \leq \epsilon 2^{l+1}$ other sets in Z' .) On

the other hand, every new member of Y' is joined to all of $Z' \cap \mathcal{P}W_2$, which has size at least $|Z \cap \mathcal{P}W_2| \geq (3/2 - 8\epsilon)2^l$. It follows that

$$e(B_2) \geq e(B_1) + C(\frac{1}{2} - 10\epsilon)2^l \geq (1 - 3\epsilon)2^{2l+1} + C(\frac{1}{2} - 10\epsilon)2^l. \quad (25)$$

We now show that $e(B_2) \leq (1 + \epsilon)2^{2l+1}$. If $|Y'| \geq 2^l$, then write $|Y'| = (1 + \phi)2^l$ where $\phi \geq 0$; Y' contains all of $\mathcal{P}W_1$, and $\phi 2^l$ ‘extra’ sets. We have $|Z'| \leq (2 - \phi)2^l$, and therefore by (23), $\phi \leq 1/2 + 6\epsilon < 1$. Note that every ‘extra’ set in $Y' \setminus \mathcal{P}W_1$ has at most 2^l neighbors in $\mathcal{P}W_2$, and therefore at most $(1 + 2\epsilon)2^l$ neighbours in Z' . Hence,

$$e(B_2) \leq 2^l(2 - \phi)2^l + \phi 2^l(1 + 2\epsilon)2^l = (1 + \phi\epsilon)2^{2l+1} \leq (1 + \epsilon)2^{2l+1}.$$

If, on the other hand, $|Y'| \leq 2^l$, then since $|Y'| + |Z'| \leq 3 \cdot 2^l$, we have $e(B_2) \leq |Y'||Z'| \leq 2^{2l+1}$. Hence, we always have

$$e(B_2) \leq (1 + \epsilon)2^{2l+1}. \quad (26)$$

Combining (25) and (26), we see that

$$C \leq \frac{8\epsilon}{1/2 - 10\epsilon}2^l \leq 20\epsilon 2^l,$$

provided $\epsilon \leq 1/100$.

This implies (a). Indeed, if $|\mathcal{P}W_1 \setminus Y| \leq C \leq 20\epsilon 2^l$, then we are done. Otherwise, by the definition of C , we have $|Y \setminus \mathcal{P}W_1| \leq 20\epsilon 2^l$. Recall that by (16), $|Y| \geq (1 - 2\epsilon)2^l$, and therefore

$$|Y \cap \mathcal{P}W_1| = |Y| - |Y \setminus \mathcal{P}W_1| \geq (1 - 2\epsilon)2^l - 20\epsilon 2^l = (1 - 22\epsilon)2^l.$$

Hence,

$$|\mathcal{P}(W_1) \setminus Y| \leq 22\epsilon 2^l, \quad (27)$$

proving (a).

Since $e(B) \geq (1 - \epsilon)2^{2l+1}$, $e(B_2) \leq (1 + \epsilon)2^{2l+1}$, and $e(B_2) \geq e(B_1)$, we have

$$e(B_1) - e(B) \leq e(B_2) - e(B) \leq (1 + \epsilon)2^{2l+1} - (1 - \epsilon)2^{2l+1} = \epsilon 2^{2l+2} \quad (28)$$

We now use this to show that

$$D = \min\{|\mathcal{P}(W_2) \setminus Z|, |Z \setminus \mathcal{P}W_2|\} \leq \sqrt{\epsilon}2^l.$$

Suppose for a contradiction that $D \geq \sqrt{\epsilon}2^l$; then it is easy to see that there must exist $z \in Z \setminus \mathcal{P}W_2$ with at least

$$2|Y|/3 - 8\sqrt{\epsilon}2^l$$

neighbours in Y . Indeed, suppose that every $z \in Z \setminus \mathcal{P}W_2$ has less than $2|Y|/3 - 8\sqrt{\epsilon}2^l$ neighbors in Y . Recall that every new member of Z' has at least $(2/3 - 8\epsilon)|Y|$ neighbours in Y . Hence,

$$e(B_1) - e(B) > 8D(\sqrt{\epsilon} - \epsilon)|Y| \geq 8\sqrt{\epsilon}2^l(\sqrt{\epsilon} - \epsilon)(1 - 2\epsilon)2^l \geq \epsilon 2^{2l+1}$$

since $\epsilon < 1/16$, contradicting (28).

Hence, we may choose $z \in Z \setminus \mathcal{P}W_2$ with at least

$$2|Y|/3 - 8\sqrt{\epsilon}2^l$$

neighbours in Y . Without loss of generality, we may assume that $n \in z \cap W_1$; then none of these neighbours can contain n . Hence, Y contains at most

$$|Y|/3 + 8\sqrt{\epsilon}2^l$$

sets containing n . But by (27), Y contains at least $(1 - 44\epsilon)2^{l-1}$ of the subsets of W_1 that contain n , and therefore $|Y| \geq (3/2 - o(1))2^l$. By (23), it follows that $|Y| = (3/2 - o(1))2^l$ and $|Z| = (3/2 + o(1))2^l$, so Y contains $(1 - o(1))2^{l-1}$ sets containing n . Hence, by (18), so does Z . As in the proof of Claim 1, we obtain a contradiction. This implies that

$$D = \min\{|\mathcal{P}(W_2) \setminus Z|, |Z \setminus \mathcal{P}W_2|\} \leq \sqrt{\epsilon}2^l,$$

as desired.

This implies (b). Indeed, if $|Z \setminus \mathcal{P}W_2| \leq \sqrt{\epsilon}2^l$, then we are done. Otherwise, by the definition of D , $|\mathcal{P}(W_2) \setminus Z| \leq \sqrt{\epsilon}2^l$, and therefore

$$|Z \cap \mathcal{P}W_2| \geq (2 - \sqrt{\epsilon})2^l.$$

Since $|Z| \leq (2 + 2\epsilon)2^l$, we have

$$|Z \setminus \mathcal{P}W_2| = |Z| - |Z \cap \mathcal{P}W_2| \leq (2 + 2\epsilon)2^l - (2 - \sqrt{\epsilon})2^l = (\sqrt{\epsilon} + 2\epsilon)2^l,$$

proving (b). $\square$

We conclude by proving the following

Claim 4.

$$|\mathcal{P}(W_2) \setminus Z| \leq 4\sqrt{\epsilon}2^l.$$

Proof. Let

$$\mathcal{F}_2 = \mathcal{P}(W_2) \setminus Z$$

be the collection of sets in $\mathcal{P}W_2$ which are missing from Z , and let

$$\mathcal{E}_1 = Y \setminus \mathcal{P}W_1$$

be the set of ‘extra’ members of Y .

Since $\mathcal{G}$ is a 2-generator for X , we can express all $|\mathcal{F}_2|2^l$ sets of the form

$$w_1 \sqcup f_2 \ (w_1 \subset W_1, f_2 \in \mathcal{F}_2)$$

as a disjoint union of two sets in $\mathcal{G}$. All but at most $\epsilon 2^{2l+1}$ of these unions correspond to edges of B . Since $|Z \setminus \mathcal{P}W_2| \leq (\sqrt{\epsilon} + 2\epsilon)2^l$, there are at most $(\sqrt{\epsilon} + 2\epsilon)2^l|Y|$ edges of B meeting sets in $Z \setminus \mathcal{P}W_2$. Call these edges of B ‘bad’, and the rest of the edges of B ‘good’. Fix $f_2 \in \mathcal{F}_2$; we can express all 2^l sets of the form

$$w_1 \sqcup f_2 \ (w_1 \subset W_1)$$

as a disjoint union of two sets in $\mathcal{G}$. If $w_1 \sqcup f_2$ is represented by a good edge, then we may write

$$w_1 \sqcup f_2 = y_1 \sqcup w_2$$

where $y_1 \in \mathcal{E}_1$ with $y_1 \cap W_1 = w_1$, and $w_2 \subset W_2$, so for every such w_1 , there is a different $y_1 \in \mathcal{E}_1$. By (24), $|Y| \leq (3/2 + 6\epsilon)2^l$, and by (27), $|Y \cap \mathcal{P}W_1| \geq (1 - 22\epsilon)2^l$, so

$$|\mathcal{E}_1| = |Y| - |\mathcal{P}(W_1) \cap Y| \leq (3/2 + 6\epsilon)2^l - (1 - 22\epsilon)2^l = (1/2 + 28\epsilon)2^l.$$

Thus, for any $f_2 \in \mathcal{F}_2$, at most $(1/2 + 28\epsilon)2^l$ unions of the form $w_1 \sqcup f_2$ correspond to good edges of B . All the other unions are generated by bad edges of B or are not generated by B at all, so

$$(1/2 - 28\epsilon)2^l|\mathcal{F}_2| \leq (2\epsilon + \sqrt{\epsilon})2^l|Y| + \epsilon 2^{2l+1}.$$

Since $|Y| \leq (3/2 + 6\epsilon)2^l$ and ϵ is small, $|\mathcal{F}_2| \leq 4\sqrt{\epsilon}2^l$, as required. $\square$

We now know that Y contains all but at most $o(2^l)$ of $\mathcal{P}W_1$, and Z contains all but at most $o(2^l)$ of $\mathcal{P}W_2$. Since $|Y| + |Z| < 3 \cdot 2^l$, we may conclude that $|Y| = (1 - o(1))2^l$ and $|Z| = (2 - o(1))2^l$. It follows from Proposition 16 that provided n is sufficiently large, we must have $\mathcal{G} = \mathcal{P}(W_1) \cup \mathcal{P}(W_2) \setminus \{\emptyset\}$, completing the proof of Theorem 4. $\square$

4 Conclusion

We have been unable to prove Conjecture 1 for $k \geq 3$ and *all* sufficiently large n . Recall that if $\mathcal{G}$ is a k -generator for an n -element set X , then

$$|\mathcal{G}| \geq 2^{n/k}.$$

In view of Proposition 18, it is natural to ask whether for any fixed k , all induced subgraphs of the Kneser graph H with $\Omega(2^{n/k})$ vertices can be made k -partite by removing at most $o(2^{2n/k})$ edges. This is false for $k = 3$, however, as the following example shows. Let n be a multiple of 6, and take an equipartition of $[n]$ into 6 sets $T_1, \dots, T_6$ of size $n/6$. Let

$$\mathcal{A} = \bigcup_{\{i,j\} \in [6]^{(2)}} (T_i \cup T_j);$$

then $|\mathcal{A}| = 15(2^{n/3})$, and $H[\mathcal{A}]$ contains a $2^{n/3}$ -blow-up of the Kneser graph $K(6, 2)$, which has chromatic number 4. It is easy to see that $H[\mathcal{A}]$ requires the removal of at least $2^{2n/3}$ edges to make it tripartite. Hence, a different argument to that in Section 3 will be required.

We believe Conjecture 1 to be true for all n and k , but it would seem that different techniques will be required to prove this.

References

- [1] Alon, N., Frankl, P., The maximum number of disjoint pairs in a family of subsets, *Graphs and Combinatorics* 1 (1985), pp. 13–21.
- [2] Alon, N., Krivelevich, M., Testing k -colorability, *SIAM J. Discrete Math.* 15 (2002), Issue 2, pp. 211–227.
- [3] Bollobás, B., Erdős, P., Simonovits, M., Szemerédi, E., Extremal graphs without large forbidden subgraphs, *Annals of Discrete Mathematics* 3 (1978) pp. 29–41.
- [4] Chung, F.K.R., Frankl, P., Graham, R.L., Shearer, J.B., Some intersection theorems for ordered sets and graphs, *J. Combin. Theory, Ser. A* 43 (1986), pp. 23–37.
- [5] Ellis, D., Note on generating all subsets of a finite set with disjoint unions, *Electron. J. Combin.* 16, (2009) no 1., Note 16.
- [6] Erdős, P., On extremal problems of graphs and generalized graphs, *Israel J. Math.* 2 (1964), pp. 183–190.
- [7] Erdős, P., On the number of complete subgraphs contained in certain graphs, *Publ. Math. Inst. Hung. Acad. Sci., Ser. A* 7 (1962), pp. 459–464.

- [8] Erdős, P., Frankl, P., Rödl, V., The asymptotic number of graphs not containing a fixed subgraph and a problem for hypergraphs having no exponent, *Graphs and Combinatorics* 2 (1986), pp. 113–121.
- [9] Erdős, P., Simonovits, M., Supersaturated graphs and hypergraphs, *Combinatorica* **3** (1983), 181–192.
- [10] Frein, Y., Lévêque, B., Sebő, A., Generating all sets with bounded unions, *Combin. Probab. Comput.* 17 (2008) pp. 641–660.
- [11] Füredi, Z., Katona, G.O.H., 2-bases of quadruples, *Combin. Probab. Comput.* 15 (2006) pp. 131–141.
- [12] Goldreich, O., Goldwasser, S., and Ron, D., Property testing and its connection to learning and approximation, *Proc. 37th Annual IEEE FOCS* (1996), pp. 339–348.

David Ellis

`D.Ellis@dpmms.cam.ac.uk`

Benny Sudakov

`b.sudakov@math.ucla.edu`